arXiv:1303.4139v1 [math.CO] 18 Mar 2013

MONOTONICITY OF THE OPTIMAL PERIMETER IN ISOPERIMETRIC PROBLEMS ON $\mathbb{Z}^k \times \mathbb{N}^d$

EMMANUEL TSUKERMAN

Abstract. In this short note, we prove general theorems for isoperimetric problems on lattices of the form $\mathbb{Z}^k \times \mathbb{N}^d$ which state that the perimeter of the optimal set is a monotonically increasing function of the volume under certain natural assumptions, such as homogeneity or local symmetry. The proved monotonicity property is surprising in view that solutions are not always nested and, consequently, standard techniques such as compressions do not apply. The monotonicity results of this note apply in particular to vertex- and edge-isoperimetric problems in the ℓ_p distances and can be used as a tool to elucidate properties of optimal sets.

1. Background

Isoperimetric problems are classical objects of study in mathematics. In general, such problems ask for sets whose boundary is smallest for a given volume. A classic example dating back to ancient Greece is to determine the shape in the plane for which the perimeter is minimized subject to a volume constraint.

Two notions of the boundary which are commonly considered are the vertex boundary and the edge boundary. These are defined as follows. Let A be a set of vertices in a graph $G = (V, E)$. The vertex boundary of A is defined to be the set $\{v \in V \setminus A : v \sim a \text{ for some } a \in A\}$. In contrast, the *edge boundary* of A is defined to be the set $\{e \in E : e = \{v, a\} \text{ for some } a \in A \text{ and } v \in V \setminus A\}$. The vertex (edge) isoperimetric problem for G asks for the minimum possible cardinality of the vertex (edge) boundary for a k-element subset of V for each $k \in \mathbb{N}$.

One of the setting in which discrete vertex- and edge-isoperimetric problems have previously been studied is the graph of the infinite grid viewed in the ℓ_1 metric, in which two nodes are adjacent whenever their distance in the ℓ_1 metric is 1 (see [1] for the vertex-isoperimetric problem and [2] for the edge-isoperimetric problem). Natural analogues of these problems are the vertex- and edge-isoperimetric problems on the infinite grid in the ℓ_∞ metric. Since the metrics ℓ_1 and ℓ_∞ are dual, one hopes for interesting connections between the two problems. Only very recently has this family of graphs began to be studied in [3], in which the vertex-isoperimetric problem is solved.

2. Monotonicity of the optimal perimeter in isoperimetric problems on $\mathbb{Z}^k \times \mathbb{N}^d$

We denote vertices of a graph G by $V(G)$, its edges by $E(G)$, and write $G = (V(G), E(G))$. For $x \in V(G)$, we set $N_{V(G)}(x) = \{y \in V(G) : \{x, y\} \in E(G)\}$ to be the vertex-neighbourhood of x and $N_{E(G)}(x) = \{e \in E(G) : e = \{x, y\} \in E(G) \text{ for some } y \in V(G)\}$ to be its edge neighbourhood. We write $N_G(x)$ for a neighbourhood of x in G when the distinction between vertex and edge does not matter.

We will be needing the following definitions:

Definition 1. *A graph* $A = (\mathbb{Z}^{k+d}, E(A))$ *is locally symmetric if for every* $x \in V(A)$*, the neighbourhood* $N_A(x)$ *of* x *is symmetric about* x*.*
A graph G *with* $V(G) = \mathbb{Z}^k \times \mathbb{N}^d$ *is locally symmetric if there exists a graph* $A = (\mathbb{Z}^{k+d}, E(A))$ *with centrally symmetric neighbourhoods such that for every* $x \in V(G)$*, the neighbourhood* $N_G(x)$ *of* x *in* G *is the intersection of the neighbourhood* $N_A(x)$ *of* x *with* G*:*

$$N_G(x) = N_A(x) \cap G.$$

If we consider the realization of an edge of a graph as a line segment connecting the two incident vertices, it is not hard to see that definition 1 implies that the the edges join to form a union of lines, each passing through at least two lattice points.

Definition 2. *A graph* $A = (\mathbb{Z}^{k+d}, E(A))$ *is homogenous if for every* $x, y \in V(A)$*, the translation sending* x *to* y *sends* $N_A(x)$ *to* $N_A(y)$*.*
A graph G *with* $V(G) = \mathbb{Z}^k \times \mathbb{N}^d$ *is homogenous if there exists a homogenous graph* $A = (\mathbb{Z}^{k+d}, E(A))$ *such that for every* $x \in V(G)$*,*

$$N_G(x) = N_A(x) \cap G.$$

Realizing edges as line segments connecting their incident vertices, one can easily verify that homogeneity implies that the line segments join to form a union of maximal sets of parallel lines. Homogeneity implies local symmetry and is a strictly stronger condition, as can be seen by the graph constructed by forming a single line by joining edges.

Definition 3. *A graph* G *is induced by a p-norm,* $0 < p \leq \infty$*, if there exists some constant* c *such that for every* $x \in V(g)$*,* $N_V(x) = \{y \in V(G) : 0 < \|x - y\|_p \leq c\}$*.*

Note that if a graph G with $V(G) = \mathbb{Z}^k \times \mathbb{N}^d$ is induced by a p-norm, then it is homogenous.

Recall that colexicographical ordering is defined by

$$(a_1, a_2, \ldots, a_n) < (b_1, b_2, \ldots, b_n) \iff (\exists m > 0)(\forall i > m)(a_i = b_i) \wedge (a_m < b_m)$$

Theorem 1. *Let* G *be a locally symmetric graph on* $\mathbb{Z}^k$*. The minimum edge-perimeter is a monotonic increasing function of the volume.*

Proof. Let B be a set of cardinality $|A| + 1$ with optimal perimeter. We find a point which can be removed without increasing the perimeter. This implies that a set of cardinality $|A|$ has minimum perimeter $|\partial A|$ less than or equal to $|\partial B|$. Such a point has the property that it has at least as many neighbours in $\mathbb{Z}^k \setminus B$ as it does in B. We derive a contradiction by assuming that such a point does not exist.

Let

$$R_x(y) = (2x_1 - y_1, 2x_2 - y_2, ..., 2x_k - y_k)$$

denote the operation of reflecting point y in point x.

Let $p \in H_r \cap B$ be greatest in colexicographical order. If p has no neighbour in B, then it can clearly be removed without increasing the perimeter. So let y be a neighbour of p in B. We show that the reflection $R_p(y)$ of y in p is not in B. Since p is greater than y colexicographically, $p_i = y_i \forall i > m$ and $p_m > y_m$ for some $m > 0$. For each $i > m$, $2p_i - y_i = p_i$. Additionally,

$$2p_m - y_m > p_m,$$

so that $R_p(y) > p$ colexicographically. Therefore $R_p(y) \notin B$. ☐

Theorem 2. *Let G be a homogenous graph on $\mathbb{Z}^k$. The minimum vertex-boundary is a monotonic increasing function of volume.*

Proof. We think of all edges as line segments connecting the vertices. Let $e = \{x, y\} \in E(G)$ be an edge of longest length. Let $\vec{e}$ be the vector having the magnitude and direction corresponding to e. Consider the family F of hyperplanes having $\vec{e}$ as a vector normal to their surface. Let $h \in F$ be the hyperplane with largest distance from the origin satisfying $h \cap B \neq \emptyset$. Take any $p \in h \cap B$. Then the point $p + \vec{e} \notin B$ and also has no neighbours in B other than p. □

Theorem 3. *Let G be a graph on $\mathbb{Z}^k \times \mathbb{N}^d$ induced by a p-norm. The minimum edge-perimeter is a monotonic increasing function of the volume.*

Proof. Being induced by a p-norm, G is locally symmetric. Therefore the argument from Theorem 1 shows that for $p \in B$ of greatest colexicographical order, the reflection $R_p(y)$ of a neighbour $y \in B$ is outside of B. However, a new issues arises, that it might land outside of the graph. For each i such that $2p_i - y_i < 0$, we apply a reflection in the hyperplane $x_i = 0$. Since orthogonal reflections commute, the order does not matter. The colexicographical order of the image can only increase, as its entries have increased. Moreover, each entry $2p_i - y_i < 0$ maps to $y_i - 2p_i > 0$, so this image is outside of B and inside the graph. Since a product of reflections is an involution, this gives an injective map from neighbours of p to points outside of B and inside the graph. Let z be the image of $R_p(y)$ under these reflections. It remains to see that z is a neighbour of p. For each i for which $2p_i - y_i < 0$, the absolute valued difference between the ith coordinates of z and p is smaller than the corresponding difference of p and y:

$$|z_i - p_i| = |(y_i - 2p_i) - p_i| \leq |y_i - p_i|.$$

For the remaining coordinates, the difference is the same. Therefore $\|z - p\|_p \geq \|y - p\|_p$, so z is a neighbour of p. □

Theorem 4. *Let G be a homogenous graph on $\mathbb{Z}^k \times \mathbb{N}^d$. The minimum vertex-perimeter is a monotonic increasing function of the volume.*

Proof. As in the proof of Theorem 2, we consider an edge $e = \{x, y\} \in E(G)$ of longest length and the family F of hyperplanes having $\vec{e}$ as a vector normal to their surface. Though point $p + \vec{e} \notin B$, this point might also be outside of the graph. If this is the case, we consider instead the vector $-\vec{e}$, which exists by the local symmetry of the graph, and proceed by the same argument. □

Finally, we note that optimal sets are not necessarily nested. Consider the graph $\mathbb{N}^n$ induced by the ∞-norm with $c = 1$. That is, the graph for which $x \sim y$ iff $\max_i |x_i - y_i| \leq 1$. The set of edges is $E(G) = \{\{x, y\} : x, y \in \mathbb{N}^n \text{ and } \|x - y\|_\infty = \max_i |x_i - y_i| = 1\}$. We consider the edge-isoperimetric problem on this graph.

Proposition 4. *The optimal sets of $\mathbb{N}^2$ are not nested.*

Proof. Figure 1 shows the uniquely determined up to reflection in $y = x$ sequence of nested optimal sets of $\mathbb{N}^2$. Figure 2 shows a set with $|A| = 11$ which has a smaller perimeter than the optimal nested set with $|A| = 11$. □

Nested optimal sets are crucial for the technique of compression and the fact that the optimal sets are not nested means that such an approach will not be possible. However, we



Figure 1. The unique (up to reflection in the line $y = x$) sequence of optimal nested sets for $1 \leq |A| \leq 11$. The optimal nested set with $|A| = 11$ has $|\partial A| = 17$.



Figure 2. An optimal set having $|A| = 11$ has $|\partial A| = 16$. This set is better than the one of the same volume in Figure 1

believe that the monotonicity of the optimal perimeter established in this note is a useful tool for obtaining bounds, proving optimality and understanding properties of optimal sets.

3. ACKNOWLEDGEMENTS

I would like to thank Professor Simon Brendle for his helpful suggestions throughout the writing of this note.

REFERENCES

[1] Wang, D.-L. and Wang, P. (1977) Discrete isoperimetric problems. *SIAM J. Appl. Math.* 32 860-870.
[2] Béla Bollobás and Imre Leader. Edge-isoperimetric inequalities in the grid. *Combinatorica*, 11(4):299-314, 1991.
[3] E. Veomett, A. J. Radcliffe.Vertex Isoperimetric Inequalities for a Family of Graphs on $\mathbb{Z}^k$. The Electronic Journal of Combinatorics **19(2)** (2012), #P45.

STANFORD UNIVERSITY, STANFORD, CA 94305
E-mail address: `emantsuk@stanford.edu`